arXiv:1701.02583v1 [physics.class-ph] 26 Dec 2016

ELECTROMAGNETIC WAVE PROPAGATION IN GLHUA INVISIBLE SPHERE BY GL NO SCATERING FULL WAVE MODELING AND INVERSION

Jianhua Li* and Feng Xie, Lee Xie, Ganquan Xie
GL Geophysical Laboratory, USA

Ganquan Xie
Chinese Dayuling Supercomputational Sciences Center, China

(Dated: December 26,2016)

Using GL no scattering full wave modeling and inversion, we create a GLHUA pre cloak electromagnetic (EM) material in the virtual sphere that makes the sphere is invisible. The invisible sphere is called GLHUA sphere. In GLHUA sphere, the Pre cloak relative parameter is not less than 1; the parameters and their derivative are continuous across the boundary r=R2 and the parameters are going to infinity at origin r=0. The phase velocity of EM wave in the sphere is less than light speed and going to zero at origin. The EM wave field excited in the outside of the sphere can not be disturbed by GLHUA sphere. By GL full wave method, we rigorously proved the incident EM wave field excited in outside of GLHUA sphere and propagation through the sphere without any scattering by the sphere, the total EM field in outside of the sphere equal to the incident wave field. Moreover, we prove that in GLHUA sphere with the pre cloak material, when r is going to origin, EM wave field propagation in GLHUA sphere is going to zero. All copyright and patent of the GLHUA EM cloaks,GLHUA sphere and GL modeling and inversion methods are reserved by authors in GL Geophysical Laboratory.

PACS numbers: 13.40.-f, 41.20.-q, 41.20.jb,42.25.Bs

I. INTRODUCTION

Using GL no scattering modeling and inversion and many GL method simulation [1] [2] [3], we find a class nonzero relative anisotropic parameter solution of EM zero scattering inversion in the sphere $r \leq R_2$ We create a novel material in GLHUA sphere with relative EM parameter not less than 1 that makes the sphere is invisible. The parameters and their derivative are continuous across the boundary $r = R_2$ and the parameters are going to infinity at origin $r = 0$. The phase velocity of EM wave in the sphere is less than light speed and going to zero at origin. We discovered and proved an essential property that in the local sphere $r \leq R_O$ without EM source, including $R_2 < r < R_O$ annular layer in free space and $r \leq R_2$ in GLHUA sphere, on any spherical surface with radius $r < R_O$, the spherical surface integral of $E_r \sin\theta$ and $H_r \sin\theta$ is zero, (E_r is radial electric wave field, H_r is radial magnetic wave field). The essential property of EM wave is a key difference from the acoustic wave and seismic wave. Based on the essential property, by GL full wave method, we rigorously proved the incident EM wave field excited in the outside of GLHUA sphere and propagation through the sphere without any scattering from GLHUA sphere, the total EM field in the outside of GLHUA sphere equal to the incident wave field. EM wave field excited in the outside of the sphere can not be disturbed by GLHUA sphere. GLHUA sphere is complete invisible. Moreover, based on the above essential property of EM wave, we prove that in GLHUA sphere with the pre cloak material, when r is going to origin, EM wave field propagation in GLHUA sphere is going to zero. We propose a special GL transform to map GLHUA outer annular layer cloak to GLHUA sphere. The anisotropic relative parameters of EM material in physical GLHUA outer annular layer cloak, $\varepsilon_{p,r} = \mu_{p,r} = 1$, $\varepsilon_{p,\theta} = \frac{1}{2}\left(\frac{r-R_1}{R_2-R_1} + \frac{R_2-R_1}{r-R_1}\right)$, $\varepsilon_{p,\theta} = \varepsilon_{p,\phi} = \mu_{p,\theta} = \mu_{p,\phi}$ are mapping to anisotropic relative parameter material in GLHUA invisible sphere that satisfy GLHUA pre invisible cloak material conditions. From the theoretical proof in GLHUA invisible sphere, we can rigorously prove GLHUA outer annular layer cloak is invisible cloak with concealment. The major new ingredients in this paper are proposed in 7 sections.The introduction is presented in the section 1. Second order Maxwell electromagnetic equation in anisotropic material in spherical coordinate, and the basic fundamental electromagnetic wave in free space is presented in the section 2. The Global and Local (GL) method for radial electromagnetic equation and the essential property of radial EM wave are presented in the section 3. In the section 4, we propose GL EM wave Greens equation and GL EM Greens function. GL EM Integral equation and its theoretical proof are proposed in the section 5. In the section 6, we propose GLHUA EM invisible sphere, in this section, we rigorously proved that EM wave field propagation is going to zero at origin in GLHUA sphere. In section 7, the discussion and conclusion is presented.

*Also at GL Geophysical Laboratory, USA, glhua@glgeo.com

II. SECOND ORDER MAXWELL ELECTROMAGNETIC EQUATION IN SPHERICAL COORDINATE

A. Electromagnetic Maxwell Equation in Sphere Coordinate System

$$\begin{aligned}\nabla\times\vec{E} &= \frac{1}{r^2\sin\theta}\begin{bmatrix}\vec{r} & r\vec{\theta} & r\sin\theta\vec{\phi}\\ \frac{\partial}{\partial r} & \frac{\partial}{\partial\theta} & \frac{\partial}{\partial\phi}\\ E_r & rE_\theta & r\sin\theta E_\phi\end{bmatrix}\\ &= -i\omega\left(H_r,H_\theta,H_\phi\right)\begin{bmatrix}\mu_r & & \\ & \mu_\theta & \\ & & \mu_\phi\end{bmatrix}\mu_0\end{aligned} \tag{1}$$

$$\begin{aligned}\nabla\times\vec{H} &= \frac{1}{r^2\sin\theta}\begin{bmatrix}\vec{r} & r\vec{\theta} & r\sin\theta\vec{\phi}\\ \frac{\partial}{\partial r} & \frac{\partial}{\partial\theta} & \frac{\partial}{\partial\phi}\\ H_r & rH_\theta & r\sin\theta H_\phi\end{bmatrix}\\ &= i\omega\left(E_r,E_\theta,E_\phi\right)\begin{bmatrix}\varepsilon_r & & \\ & \varepsilon_\theta & \\ & & \varepsilon_\phi\end{bmatrix}\varepsilon_0+\delta(\vec{r}-\vec{r}_s)\vec{e}_s\end{aligned} \tag{2}$$

$$\begin{aligned}&\nabla\cdot\vec{B} = \tfrac{1}{r^2}\tfrac{\partial}{\partial r}\left(r^2\mu_r H_r\right)+\tfrac{1}{\sin\theta r}\tfrac{\partial}{\partial\theta}\sin\theta\mu_\theta H_\theta\\ &+\tfrac{1}{r\sin\theta}\tfrac{\partial}{\partial\phi}\mu_\phi H_\phi = 0,\end{aligned} \tag{3}$$

$$\begin{aligned}&\nabla\cdot\vec{D} = \tfrac{1}{r^2}\tfrac{\partial}{\partial r}\left(r^2\varepsilon_r E_r\right)+\tfrac{1}{\sin\theta r}\tfrac{\partial}{\partial\theta}\sin\theta\varepsilon_\theta E_\theta\\ &+\tfrac{1}{r\sin\theta}\tfrac{\partial}{\partial\phi}\varepsilon_\phi E_\phi = 4\pi\rho,\end{aligned} \tag{4}$$

where ρ is the electric charge, ε_0 is the basic constant electric permittivity, μ_0 is the basic constant magnetic permeability, ε_r , relative radial electric permittivity, ε_θ and ε_ϕ, relative angular electric permittivity, μ_r , relative radial magnetic permeability, μ_θ and μ_ϕ, relative angular magnetic permeability, E_r is the radial electric field, E_θ is the electric field in angular θ direction, E_ϕ is the electric field in angular ϕ direction, $\vec{E}=(E_r,E_\theta,E_\phi)$ is the electric field vector, $\vec{D}$ is the electric displacement vector, $\vec{B}$ is the magnetic flux vector, H_r is the radial magnetic field, H_θ is the magnetic field in angular θ direction, H_ϕ is the magnetic field in angular ϕ direction, $\vec{H}=(H_r,H_\theta,H_\phi)$ is the magnetic field vector, $\omega=2\pi f$ is the angular frequency, f is the frequency Hz, $\vec{r}=(r,\theta,\phi)$, the vector in spherical coordinate.

B. Second Order Maxwell Electromagnetic Equation On Radial Electric Wave In Spherical Coordinate

We use Electric-Magnetic-Electric EME translate process to translate Maxwell Equation in Spherical Coordinate (S1) and (S2) with anisotropic material into the Second order Maxwell Electromagnetic Equation on Radial Electric Wave Field In Spherical Coordinate

$$\begin{aligned}&\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\tfrac{\partial}{\partial r}\varepsilon_r r^2E_r+\tfrac{1}{\sin\theta}\tfrac{\partial}{\partial\theta}\sin\theta\tfrac{\partial E_r}{\partial\theta}+\\ &\tfrac{1}{\sin^2\theta}\tfrac{\partial^2E_r}{\partial\phi^2}+k^2\mu_\theta\varepsilon_r r^2E_r\\ &=-\tfrac{1}{\sin\theta}\tfrac{\partial}{\partial\theta}\sin\theta\tfrac{\partial}{\partial r}\left(\tfrac{1}{i\omega\varepsilon_\theta}rJ_\theta\right)\\ &-\tfrac{1}{\sin\theta}\tfrac{\partial}{\partial\phi}\tfrac{\partial}{\partial r}\left(\tfrac{1}{i\omega\varepsilon_\theta}rJ_\phi\right)\\ &-\tfrac{\partial}{\partial r}\tfrac{1}{i\omega\varepsilon_\theta}\tfrac{\partial}{\partial r}r^2J_r-(-i\omega\mu_\theta)r^2J_r,\end{aligned} \tag{5}$$

C. Second order Maxwell Electromagnetic Equation on Radial Magnetic Wave Field In Spherical Coordinate

We use MEM Magnetic-Electric-Magnetic translate process to translate Maxwell equation in spherical Coordinate (S1) and (S2) into the Second order Maxwell magnetic equation on radial magnetic wave field in spherical coordinate with anisotropic material,

$$\begin{aligned}&\tfrac{\partial}{\partial r}\tfrac{1}{(\mu_\theta)}\tfrac{\partial\left(\mu_r r^2H_r\right)}{\partial r}+\tfrac{1}{\sin\theta}\tfrac{\partial}{\partial\theta}\sin\theta\tfrac{\partial H_r}{\partial\theta}\\ &+\tfrac{1}{\sin^2\theta}\tfrac{\partial H_r}{\partial\phi^2}+k^2\varepsilon_\theta\mu_r r^2H_r\\ &=\tfrac{1}{\sin\theta}\tfrac{\partial}{\partial\theta}\sin\theta rJ_\phi-\tfrac{1}{\sin\theta}\tfrac{\partial}{\partial\phi}rJ_\theta,\end{aligned} \tag{6}$$

D. Second Order Maxwell Electromagnetic Equation On Angular Electric Wave Field In Spherical Coordinate

After solving equation (5) and (6) and obtaining the radial electric wave field $E_r(r,\theta,\phi)$ and radial magnetic wave field $H_r(r,\theta,\phi)$, the following second order Maxwell electromagnetic equation govern angular electric wave field $E_\theta(r,\theta,\phi)$ and $E_\phi(r,\theta,\phi)$

$$\begin{aligned}&\tfrac{\partial}{\partial r}\tfrac{1}{(\mu_\theta)}\tfrac{\partial rE_\theta}{\partial r}+k^2\varepsilon_\theta rE_\theta\\ &=-i\omega\mu_0\tfrac{1}{\sin\theta}\tfrac{\partial H_r}{\partial\phi}+\tfrac{\partial}{\partial r}\left(\tfrac{1}{\mu_\theta}\tfrac{\partial E_r}{\partial\theta}\right)\\ &-i\omega\mu_0rJ_\theta\end{aligned} \tag{7}$$

$$\begin{aligned}&\tfrac{\partial}{\partial r}\tfrac{1}{(\mu_\theta)}\tfrac{\partial rE_\phi}{\partial r}+k^2\varepsilon_\theta rE_\phi\\ &=\tfrac{\partial}{\partial r}\tfrac{1}{(\mu_\theta)}\tfrac{1}{\sin\theta}\tfrac{\partial E_r}{\partial\phi}+i\omega\mu_0\tfrac{\partial H_r}{\partial\theta}\\ &-i\omega\mu_0rJ_\phi\end{aligned} \tag{8}$$

E. Second Order Maxwell Electromagnetic Equation on Angular Magnetic Wave Field In Spherical Coordinate

After solving equation (5) and (6) and obtaining the radial electric wave field $E_r(r,\theta,\phi)$ and radial magnetic wave field $H_r(r,\theta,\phi)$, the following second order Maxwell electromagnetic equation govern angular magnetic wave field $H_\theta(r,\theta,\phi)$ and $H_\phi(r,\theta,\phi)$

$$\begin{array}{l}\frac{\partial}{\partial r}\left(\frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}rH_\theta\right)+k^2\mu_\theta rH_\theta= \\ i\omega\varepsilon_0\frac{1}{\sin\theta}\frac{\partial E_r}{\partial\phi}+\frac{\partial}{\partial r}\left(\frac{1}{\varepsilon_\theta}\left(\frac{\partial H_r}{\partial\theta}\right)\right) \\ +\frac{\partial}{\partial r}\left(\frac{1}{\varepsilon_\theta}rJ_\phi\right),\end{array} \tag{9}$$

$$\begin{array}{l}\frac{\partial}{\partial r}\left(\frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}rH_\phi\right)+k^2\mu_\theta rH_\phi= \\ \frac{1}{\sin\theta}\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\frac{\partial H_r}{\partial\phi}-i\omega\varepsilon_0\frac{\partial E_r}{\partial\theta} \\ -\frac{\partial}{\partial r}\left(\frac{1}{\varepsilon_\theta}rJ_\theta\right),\end{array} \tag{10}$$

F. Fundamental acoustic wave field

Let $g(\vec{r},\vec{r}_s)$ is the fundamental acoustic wave field, which is solution of following acoustic equation

$$\begin{array}{l}\frac{\partial}{\partial r}r^2\frac{\partial g}{\partial r}+\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial g}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2 g}{\partial\phi^2} \\ +k^2r^2g=\delta(r-r_s)\frac{\delta(\theta-\theta_s)}{\sin\theta}\delta(\phi-\phi_s),\end{array} \tag{11}$$

$$g(\vec{r},\vec{r}_s)=-\frac{1}{4\pi}\frac{e^{-ik|\vec{r}-\vec{r}_s|}}{|\vec{r}-\vec{r}_s|} \tag{12}$$

$$\begin{array}{l}|\vec{r}-\vec{r}_s|=\sqrt{|\vec{r}-\vec{r}_s|^2}, \\ |\vec{r}-\vec{r}_s|^2=r^2+r_s^2 \\ -2rr_s\sin\theta\sin\theta_s\cos(\phi-\phi_s) \\ -2rr_s\cos\theta\cos\theta_s,\end{array} \tag{13}$$

where $\vec{r}_s=(r_s,\theta_s,\phi_s)$ is point source location, $\vec{r}=(r,\theta,\phi)$ is variable spherical coordinate, i.e. observation point.

G. Basic fundamental electromagnetic wave field in free space

The basic fundamental electromagnetic wave field in free space can be excited by current source $\vec{J}=\delta(\vec{r}-\vec{r}_s)\vec{e}_j$, or magnetic moment source $\vec{M}=\delta(\vec{r}-\vec{r}_s)\vec{e}_j$.We consider the current source in this paper, similarly theorem and proof are suitable for the magnetic moment source. Let $\varepsilon_r=\varepsilon_\theta=\varepsilon_\phi=1$ and $\mu_r=\mu_\theta=\mu_\phi=1$, electric current source $\vec{J}=\delta(\vec{r}-\vec{r}_s)\vec{e}_j$,The basic fundamental electromagnetic wave field in free space are solutions of equations (5)-(6) in free space,

$$E^b{}_j=\frac{1}{i\omega\varepsilon_0}\left(\nabla\nabla\cdot(g\vec{e}_j)+k^2g\vec{e}_j\right) \tag{14}$$

$$\begin{array}{l}E^b{}_{j,r}=\frac{1}{i\omega\varepsilon_0}\left(e_{jr}\left(\frac{\partial^2 g}{\partial r^2}+k^2g\right)+e_{j\theta}\frac{\partial}{\partial r}\frac{1}{r}\frac{\partial g}{\partial\theta}\right) \\ +e_{j\phi}\frac{1}{i\omega\varepsilon_0}\frac{\partial}{\partial r}\frac{1}{r\sin\theta}\frac{\partial g}{\partial\phi}\end{array} \tag{15}$$

$$\begin{array}{l}H^b{}_j=\nabla\times(g\vec{e}_j), \\ H^b{}_{j,r}=\left(e_{j\phi}\frac{1}{r}\frac{\partial g}{\partial\theta}-e_{j\theta}\frac{1}{r\sin\theta}\frac{\partial g}{\partial\phi}\right),\end{array} \tag{16}$$

where the above equation are in the spherical coordinate system, $g(\vec{r},\vec{r}_s)$ is denoted in (11)-(13), $\nabla\cdot$ is diverge operator E^b_j is basic fundamental electric wave, , H^b_j is basic fundamental magnetic wave, which is excited by source $\vec{J}=\delta(\vec{r}-\vec{r}_s)\vec{e}_j$, the up script b means basic fundamental electric wave, the lower script j means source with excited vector $\vec{e}_j$. $E^b_{j,r}$ is radial basic fundamental electric wave, i.e. r component of E^b_j , $H^b_{j,r}$ is radial basic fundamental magnetic wave, i.e. r component of H^b_j , $\vec{e}_j$, j=1,2,3, is a unite vector,

$$\begin{array}{l}e_1=e_x=\left(\begin{array}{ccc}\sin\theta\cos\phi & \cos\theta\cos\phi & -\sin\phi\end{array}\right), \\ e_2=e_y=\left(\begin{array}{ccc}\sin\theta\sin\phi & \cos\theta\sin\phi & \cos\phi\end{array}\right), \\ e_3=e_z=\left(\begin{array}{ccc}\cos\theta & -\sin\theta & 0\end{array}\right),\end{array} \tag{17}$$

III. GLOBAL AND LOCAL (GL) METHOD FOR RADIAL ELECTROMAGNETIC EQUATION AND THE ESSENTIAL PROPERTY OF THE RADIAL ELECTROMAGNETIC WAVE

In the section 1, we proposed second order Maxwell electromagnetic equation in anisotropic material in spherical coordinate. The fundamental electromagnetic wave field in free space in sphere coordinate is presented in section 2. These basic equations are used in this and next sections.

A. Global and Local GL radial electric second order differential equation

For $R_2>0$, we consider electromagnetic equation (5)-(10) in the sphere $r\le R_2$ with anisotropic media, in the outside of the sphere, $r>R_2$ with basic isotropic electric permittivity ε_0 and magnetic permeability μ_0 in free space. In our paper, we suppose that the electromagnetic source set is bounded, the bounded source set Ω_s is in outside of the large sphere with radius R_O, $R_O>R_2$,for example, the point source located in outside of the sphere, $r_s>R_O>R_2$.

Using Global and Local (GL) field method, we propose radial GL electromagnetic field and GL electromagnetic equation. and study the following GL radial electromagnetic equation in the sphere$r\le R_2$.

Definition of GL radial electromagnetic wave field

$$\begin{array}{l}E(\vec{r})=\varepsilon_r r^2E_r(\vec{r}), \\ H(\vec{r})=\mu_r r^2H_r(\vec{r}),\end{array} \tag{18}$$

From the radial electric equation (5) , we propose GL radial electric second order differential equation

$$\begin{array}{l}\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}E+\frac{1}{\varepsilon_r r^2}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial E}{\partial\theta} \\ +\frac{1}{\varepsilon_r r^2}\frac{1}{\sin^2\theta}\frac{\partial^2 E}{\partial\phi^2}+k^2\mu_\theta E=J_s,\end{array} \tag{19}$$

where $k = 2\pi f\sqrt{\varepsilon_0\mu_0}$,

$$J_s = \delta(r-r_s)\tfrac{1}{\sin\theta}\delta(\theta-\theta_s)\delta(\phi-\phi_s)e_r$$

electric point source $r_s > R_O$. The incident GL electric wave field in free space,

$$\begin{array}{l} E_j^b(\vec{r}) = r^2 E_{j,r}^b = \frac{1}{i\omega\varepsilon_0} r^2 e_{jr}\left(\frac{\partial^2 g}{\partial r^2} + k^2 g\right) \\ + \frac{1}{i\omega\varepsilon_0} r^2\left(e_{j\theta}\frac{\partial}{\partial r}\frac{1}{r}\frac{\partial g}{\partial\theta} + e_{j\phi}\frac{\partial}{\partial r}\frac{1}{r\sin\theta}\frac{\partial g}{\partial\phi}\right), \end{array} \tag{20}$$

where $g = g(r, r_s)$ is the fundamental solution of the acoustic wave equation, (11-13). Let $E^b(\vec{r})$ to denote one of $E_j^b(\vec{r}) = r^2 E_{j,r}^b$, $j = 1, 2, 3$, it obvious that

$$\lim_{r\to 0} E^b(\vec{r}) = 0, \tag{21}$$

$$\lim_{r\to 0}\frac{\partial}{\partial r}E^b(\vec{r}) = 0, \tag{22}$$

B. Global and Local GL radial magnetic field second order differential equation

By definition of GL magnetic wave $H(\vec{r}) = \mu_r r^2 H_r(\vec{r})$ in (18), from the radial magnetic equation (6), we propose GL radial magnetic field second order differential equation

$$\begin{array}{l} \frac{\partial}{\partial r}\frac{1}{\mu_\theta}\frac{\partial}{\partial r}H + \frac{1}{\mu_r r^2}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial H}{\partial\theta} \\ + \frac{1}{\mu_r r^2}\frac{1}{\sin^2\theta}\frac{\partial^2 H}{\partial\phi^2} + k^2\varepsilon_\theta H = M_s, \end{array} \tag{23}$$

M_s is magnetic point source, incident GL magnetic wave in free space is,

$$\begin{array}{l} H_j^b = r^2 H_{j,r}^b \\ = r\left(e_{j\phi}\frac{\partial}{\partial\theta}g - e_{j\theta}\frac{1}{\sin\theta}\frac{\partial}{\partial\phi}g\right), \end{array} \tag{24}$$

Let $H^b(\vec{r})$ to denote one of the $H_j^b(\vec{r}) = r^2 H_{j,r}^b$, $j = 1, 2, 3$, it obvious

$$\lim_{r\to 0} H^b(\vec{r}) = 0, \tag{25}$$

$$\lim_{r\to 0}\frac{\partial}{\partial r}H^b(\vec{r}) = 0, \tag{26}$$

C. Spherical surface integral of incident GL electromagnetic wave is vanished

Define spherical surface integral of incident GL electric wave in free space as

$$E_0^b(r) = \frac{1}{4\pi}\int_0^\pi\int_0^{2\pi} E^b(\vec{r})\sin\theta d\theta d\phi, \tag{27}$$

Define spherical surface integral of incident GL magnetic wave in free space as

$$H_0^b(r) = \frac{1}{4\pi}\int_0^\pi\int_0^{2\pi} H^b(\vec{r})\sin\theta d\theta d\phi, \tag{28}$$

D. Essential property of GL radial electromagnetic wave

***Theorem* 3.1 :**, Suppose that the electromagnetic source set is bounded, the bounded source set is in outside of the sphere with large radius R_O, $r_s > R_O$. In the no source domain, with weight $\sin\theta$ spherical surface integral of incident radial GL electromagnetic wave is zero.

$$E_0^b(r) = 0, H_0^b(r) = 0, \tag{29}$$

The spherical surface integral of radial GL electromagnetic wave is zero,

$$E_0(r) = 0, H_0(r) = 0, \tag{30}$$

***Proof* :**, By equation (19), the GL radial electric second order differential equation in free space is

$$\begin{array}{l} \frac{\partial^2}{\partial r^2}E^b + \frac{1}{r^2}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial E^b}{\partial\theta} \\ + \frac{1}{r^2}\frac{1}{\sin^2\theta}\frac{\partial^2 E^b}{\partial\phi^2} + k^2 E^b = S_E \end{array} \tag{31}$$

The electromagnetic source S_E is denoted by (5), (6), because the bounded source set is in outside of the sphere, with large radius R_O,$r_s > R_O$, there exist the no source domain $r < R_O$, in the no source domain or for plane electromagnetic wave without source,$S_E = 0$. Use $\sin\theta$ times both sides of (31) and take spherical surface integral and by integral by parts, we get Linville ordinary equation

$$\frac{d^2}{dr^2}E_0^b + k^2 E_0^b = 0, \tag{32}$$

From (21) and (22), the initial condition is

$$\lim_{r\to 0} E_0^b(r) = 0 \tag{33}$$

$$\lim_{r\to 0}\frac{\partial}{\partial r}E_0^b(r) = 0, \tag{34}$$

The equation system (32)-(34) has only zero solution. We have proved that $E_0^b(r)$ is complete vanished

$$E_0^b(r) = 0,$$

Similarly, we have proved that $H_0^b(r)$ is complete vanished,

$$H_0^b(r) = 0.$$

The first part (29) of the theorem 3.1 is proved. Next, we prove the second part of the theorem. Suppose that the electromagnetic source set is bounded, the bounded source set is in outside of the sphere with radius R_O, in the GLHUA sphere, $r \leq R_2 < R_O$, the source term is zero in the right hand of GL electric equation (19). Use $\sin\theta$ times both sides of (19) and take sphere surface integral and by integral by parts, we have ordinary equation

$$\begin{array}{l} \frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}E_0(r) + k^2\mu_\theta E_0(r) = 0, \\ r \leq R_2, \end{array} \tag{35}$$

By above proof, we have initial condition $E_0^b(R_2) = 0$ and $\frac{\partial}{\partial r}E_0^b(R_2) = 0$. Because in the GLHUA sphere, electromagnetic material parameters and their derivative are continuous across the outer boundary $r = R_2$, the radial GL electromagnetic wave and their derivative are continuous across the boundary $r = R_2$.

$$E_0(R_2) = E_0^b(R_2) = 0, \tag{36}$$

$$\frac{\partial}{\partial r}E_0(R_2) = \frac{\partial}{\partial r}E_0^b(R_2) = 0, \tag{37}$$

The solution of ordinary differential equation (35) with zero initial boundary condition (36) and (37) must be zero. Therefore,$E_0(r) = 0$ similar $H_0(r) = 0$, we proved theorem 3.1 that spherical surface integral of GL electromagnetic wave is vanished.

***Theorem* 3.2 :** Spherical surface integral of incident radial electromagnetic wave is zero. Spherical surface integral of radial electromagnetic wave is zero.

***Proof* :**,

$$\begin{array}{l} E_{0r}^b(r) = \\ \frac{1}{4\pi r^2}\int_0^\pi \int_0^{2\pi} E^b(\vec{r})\sin\theta d\theta d\phi = 0, \end{array} \tag{38}$$

$$\begin{array}{l} H_{0r}^b(r) = \\ = \frac{1}{4\pi r^2}\int_0^\pi \int_0^{2\pi} H^b(\vec{r})\sin\theta d\theta d\phi = 0, \end{array} \tag{39}$$

$$\begin{array}{l} E_{0,r}(r) = \\ = \frac{1}{4\pi r^2}\int_0^\pi \int_0^{2\pi} E(\vec{r})\sin\theta d\theta d\phi = 0, \end{array} \tag{40}$$

$$\begin{array}{l} H_{0,r}(r) = \\ = \frac{1}{4\pi r^2}\int_0^\pi \int_0^{2\pi} H(\vec{r})\sin\theta d\theta d\phi = 0 \end{array} \tag{41}$$

For EM point source and $r_s > R_O$, and $r < R_O$,also from (15) and (16), by direct integral, we can calculate

$$\begin{array}{l} E_{0jr}^b(r) = \\ = \frac{1}{4\pi}\int_0^\pi \int_0^{2\pi} E_{jr}^b(\vec{r})\sin\theta d\theta d\phi = 0, \end{array} \tag{42}$$

and

$$\begin{array}{l} H_{0jr}^b(r) = \\ = \frac{1}{4\pi}\int_0^\pi \int_0^{2\pi} H_{jr}^b(\vec{r})\sin\theta d\theta d\phi = 0, \end{array} \tag{43}$$

Here, we direct integral of (43), because (16)

$$\begin{array}{l} H_{jr}^b = \frac{1}{r^2\sin\theta}\frac{\partial}{\partial\theta}(r\sin\theta e_{j\phi}g) \\ -\frac{1}{r^2\sin\theta}\frac{\partial}{\partial\phi}(re_{j\theta}g), j = 1,2,3, \end{array}$$

$\vec{e}_j$ is an unit vector of the source, j=1,2,3, in (17),

$$\begin{array}{l} H_{0jr}^b(r) = \frac{1}{4\pi}\int_0^\pi \int_0^{2\pi} H_{jr}^b(\vec{r})\sin\theta d\theta d\phi \\ = \frac{1}{4\pi}\frac{1}{r^2}\int_0^{2\pi}\int_0^\pi \frac{\partial}{\partial\theta}(r\sin\theta e_{j\phi}g)d\theta d\phi \\ -\frac{1}{4\pi}\frac{1}{r^2}\int_0^\pi \int_0^{2\pi} \frac{\partial}{\partial\phi}(re_{j\theta}g)d\phi d\theta = 0, \end{array}$$

(43) is already proved by direct integral. Similarly, by direct integral, we can prove (42).

From theorem 3.1,$E_0(r) = 0$, by definition of GL electromagnetic wave field

$$\begin{array}{l} E(\vec{r}) = \varepsilon_r r^2 E_r(\vec{r}), \\ H(\vec{r}) = \mu_r r^2 H_r(\vec{r}), \end{array}$$

we have

$$E_{0,r}(\vec{r}) = \frac{1}{\varepsilon_r r^2}E_0(\vec{r}) = 0$$

and

$$H_{0.r}(\vec{r}) = \frac{1}{\mu_r r^2}H_0(\vec{r}) = 0$$

. The theorem 3.2 is proved.

The spherical surface integral of incident radial electromagnetic wave is zero that is essential property. The key property is essential different between electromagnetic wave and acoustic wave. Note that GL electromagnetic wave (18) is not Maxwell electromagnetic field wave, also is not flux nor displace current. In the GL method, Global and Local virtual wave E and H in (18) is convenient under any coordinate transform. Global and Local virtual wave (18) and GL second order differential equation (19) (23) are important for study GLHUA sphere and GLHUA cloak. It is shown that the GL electromagnetic differential equation and their incident wave (18)-(21) and (22)-(25) have sane equation form and theoretical properties. For simply, we study GL electric differential equation and its incident wave (18)-(21) in detail.

IV. GL ELECTROMAGNETIC GREENS EQUATION

A. We propose GL electromagnetic Greens equation for GL electric equation (18) and GL magnetic equation (23)

$$\begin{array}{l}\frac{\partial}{\partial r}\frac{\partial}{\partial r}G(\vec{r},\vec{r}')+\frac{1}{r^2}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial}{\partial\theta}G(\vec{r},\vec{r}')\\+\frac{1}{r^2}\frac{1}{\sin^2\theta}\frac{\partial^2}{\partial\phi^2}G(\vec{r},\vec{r}')\\+k^2G(\vec{r},\vec{r}')=\delta(\vec{r}-\vec{r}'),\end{array}\tag{44}$$

B. GL electromagnetic Greens function

We find and propose GL Greens function $G(\vec{r},\vec{r}')$ which is the solution of above GL electromagnetic Greens equation (44),

$$G(\vec{r},\vec{r}')=rr'g(\vec{r},\vec{r}'),\tag{45}$$

$$g(\vec{r},\vec{r}')=-\frac{1}{4\pi}\frac{e^{-ik|\vec{r}-\vec{r}'|}}{|\vec{r}-\vec{r}'|},\tag{46}$$

Where

$$\begin{array}{l}|\vec{r}-\vec{r}'|=\sqrt{|\vec{r}-\vec{r}'|^2},\\|\vec{r}-\vec{r}'|^2=r^2-2rr'\sin\theta\sin\theta'\cos(\phi-\phi')\\-2rr'\cos\theta\cos\theta'+r'^2,\end{array}$$

The GL Greens equation (44) and GL Greens function (45) are suitable for all global free space which is as background of local cloak.

C. The spherical surface integral of the GL electromagnetic Greens equation

We take

$$\begin{array}{l}G_0(r,r')=\\\frac{1}{4\pi}\int_0^{\pi}\int_0^{2\pi}G(\vec{r},\vec{r}')\sin\theta d\theta d\phi,\end{array}\tag{47}$$

Take sphere surface integral of GL electromagnetic Greens equation (44), then sphere surface integral of Greens function, $G_0(r,r')$, in (44) satisfy

$$\begin{array}{l}\frac{\partial}{\partial r}\frac{\partial}{\partial r}G_0(r,r')\\+k^2G_0(r,r')=\delta(r-r'),\end{array}\tag{48}$$

We find spherical surface integral of GL Greens function $G_0(r,r')$ which is the solution of above equation (48),

$$\begin{array}{l}G_0(r,r')=\\rr'ikj_0(kr)\left(j_0(kr')-iy_0(kr')\right),\\r\le r',\\G_0(r,r')=\\rr'ikj_0(kr')\left(j_0(kr)-iy_0(kr)\right),\\r'\le r,\end{array}\tag{49}$$

$$\begin{array}{l}j_0(kr)=\frac{\sin kr}{kr},\\y_0(kr)=-\frac{\cos kr}{kr},\end{array}\tag{50}$$

V. GL ELECTROMAGNETIC INTEGRAL EQUATION

We propose the Global and Local GL integral equation on GL electric wave field $E(\vec{r})$ in (18) in the sphere body $r\le R_2$

$$\begin{array}{l}E(\vec{r}')=E^b(\vec{r}')-\\\int\limits_{S(r\le R_2)}\left(1-\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}G\frac{\partial}{\partial r}EdV\\+\int\limits_{S(r\le R_2)}\frac{1}{r^2}\left(1-\frac{1}{\varepsilon_r}\right)\\\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2G}{\partial\phi^2}\right)EdV\\+\int\limits_{S(r\le R_2)}k^2\left(1-\mu_\theta\right)GEdV,\end{array}\tag{51}$$

The equivalent between GL integral equation (51) and GL electric wave differential equation (19) is proved next theorem 5.1. In integral equation (51), we change E to H, ε_r to μ_r, ε_θ to μ_θ, we obtain GL magnetic integral equation.

$$\begin{array}{l}H(\vec{r}')=H^b(\vec{r}')\\-\int\limits_{S(r\le R_2)}\left(1-\frac{1}{\mu_\theta}\right)\frac{\partial}{\partial r}G\frac{\partial}{\partial r}HdV\\+\int\limits_{S(r\le R_2)}\frac{1}{r^2}\left(1-\frac{1}{\mu_r}\right)\\\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2G}{\partial\phi^2}\right)HdV\\+\int\limits_{S(r\le R_2)}k^2\left(1-\varepsilon_\theta\right)GHdV,\end{array}\tag{52}$$

A. The equivalent between GL integral equation (51) and GL electric wave differential equation (19)

***Theorem* 5.1 :** Suppose that the radial electric wave $E(\vec{r})$ is solution of the GL radial electric wave differential equation (19) with incident wave (20)-(22), Greens function $G(\vec{r},\vec{r}')$ in (45) satisfy the GL electromagnetic Greens differential equation (44), then $E(\vec{r})$ satisfy the GL integral equation (51)

***Proof* :** By using Greens function $G(\vec{r},\vec{r}')$ in (45) to time the GL electric differential equation (19) and after

doing some calculation and integral by part, we have

$$\begin{array}{l}\int\limits_{S(r\le R_2)} \frac{\partial}{\partial r}\left(\frac{1}{\varepsilon_\theta}\left(\frac{\partial}{\partial r}E\right)G\right)dV \\ -\int\limits_{S(r\le R_2)} \frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}E\frac{\partial}{\partial r}GdV \\ +\int\limits_{S(r\le R_2)} \left(\frac{1}{\varepsilon_r r^2}\right. \\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2 G}{\partial\phi^2}\right)\right)EdV \\ +\int\limits_{S(r\le R_2)} k^2\mu_\theta GEdV = E^b(\vec{r})\end{array} \tag{53}$$

By using unknown wave function $E(\vec{r},\vec{r}_s)$ to time the GL Greens equation (44) and after do some calculation and integral by part, we have

$$\begin{array}{l}\int\limits_{S(r\le R_2)} \frac{\partial}{\partial r}\left(\left(\frac{\partial}{\partial r}G\right)E\right)dV \\ -\int\limits_{S(r\le R_2)} \frac{\partial}{\partial r}G\frac{\partial}{\partial r}EdV \\ +\int\limits_{S(r\le R_2)} \left(\frac{1}{r^2}\right. \\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2 G}{\partial\phi^2}\right)\right)EdV \\ +\int\limits_{R^3} k^2GEdV = E(\vec{r}),\end{array} \tag{54}$$

To subtract (53) from (54)

$$\begin{array}{l}E(\vec{r}') = E^b(\vec{r}') \\ +\frac{e^{-ikr'}}{4\pi}\int_0^\pi\int_0^{2\pi} E(0,\theta,\phi)\sin\theta d\theta d\phi \\ -\int\limits_{S(r\le R_2)} \left(1-\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}G\frac{\partial}{\partial r}EdV \\ +\int\limits_{S(r\le R_2)} \left(\frac{1}{r^2}\left(1-\frac{1}{\varepsilon_r}\right)\right. \\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2 G}{\partial\phi^2}\right)\right)EdV \\ +\int\limits_{S(r\le R_2)} k^2\left(1-\mu_\theta\right)GEdV.\end{array} \tag{55}$$

Substitute (27) for sphere surface integral of electric wave into the (55), because the anisotropic inhomogeneous electromagnetic relative material parameter are variable in the sphere,$r \le R_2$ The outside of sphere, $r > R_2$ is free space with relative electric permittivity $diag(1,1,1)$ and magnetic permeability $diag(1,1,1)$, because (30)

$$\frac{e^{-ikr'}}{4\pi}\int_0^\pi\int_0^{2\pi} E(0,\theta,\phi)\sin\theta d\theta d\phi = 0$$

the integral equation (55) will become

$$\begin{array}{l}E(\vec{r}') = E^b(\vec{r}') \\ -\int\limits_{S(r\le R_2)} \left(1-\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}G\frac{\partial}{\partial r}EdV \\ +\int\limits_{S(r\le R_2)} \left(\frac{1}{r^2}\left(1-\frac{1}{\varepsilon_r}\right)\right. \\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2 G}{\partial\phi^2}\right)\right)EdV \\ +\int\limits_{S(r\le R_2)} k^2\left(1-\mu_\theta\right)GEdV.\end{array} \tag{51}$$

The theorem 5.1 is proved.

VI. THEORY OF GLHUA ELECTROMAGNETIC INVISIBLE SPHERE AND BEHAVIOR OF THE ELECTROMAGNETIC WAVE FIELD PROPAGATION

A. GLHUA pre cloak material conditions on relative electric permittivity and magnetic permeability for invisible sphere

In this section, we propose GLHUA pre cloak material conditions of anisotropic relative electric permittivity and magnetic permeability for invisible virtual sphere $r \le R_2$, which are of the following properties for invisible sphere $r \le R_2$. The electric permittivity is the product of relative electric permittivity and ε_0, the magnetic permeability is the product of relative magnetic permeability and $\mu_0.\varepsilon_0$ Is the basic constant electric permittivity, μ_0 is basic constant magnetic permeability. GLHUA pre cloak material conditions in invisible virtual sphere $r \le R_2$ are as follows:

$$\begin{array}{l}(6.1)\mu_r(r) = \varepsilon_r(r), \\ \mu_\theta(r) = \mu_\phi(r) = \varepsilon_\theta(r) = \varepsilon_\phi(r), \\ \text{in the sphere } r \le R_2 \text{ are continuous} \\ \text{differentiable function of r,}\end{array} \tag{56}$$

$$\begin{array}{l}(6.2)\text{these parameter functions} \\ \text{and their derivative functions} \\ \text{are continuous across boundary} \\ r = R_2,\end{array} \tag{57}$$

$$\begin{array}{l}(6.3)\lim\limits_{r\to 0} r^2\mu_r(r) = \\ \lim\limits_{r\to 0} r^2\varepsilon_r(r) = \infty,\end{array} \tag{58}$$

$$\begin{array}{l}(6.4)\mu_\theta(r) = \mu_\phi(r) \\ = \varepsilon_\theta(r) = \varepsilon_\phi(r) = f(r)\frac{1}{r^2}, \\ \text{and their derivative are} \\ \text{continuous across boundary} \\ r = R_2, \lim\limits_{r\to 0} f(r) = \frac{R_2^2}{2}, \\ \lim\limits_{r\to 0}\frac{1}{r}f'(r) = 0,\end{array} \tag{59}$$

B. GL electromagnetic field and are approaching to zero at r=0 in invisible sphere

***Theorem* 6.1 :** Suppose that the anisotropic relative electric permittivity$\varepsilon_r(r)$,$\varepsilon_\theta(r)$ $\varepsilon_\phi(r)$ and magnetic permeability, $\mu_r(r)$,$\mu_\theta(r)$ $\mu_\phi(r)$, satisfy the above GL-HUA pre cloak material conditions in invisible sphere (6.1) *to* (6.4), also we suppose that

$$\int\limits_{S(r\leq R_2)} \left(|E(\vec{r})|^2+|H(\vec{r})|^2\right) dV is finite, \tag{60}$$

then GL radial electromagnetic wave field is vanished at origin,$r=0$,

$$\lim_{r\to 0} E(\vec{r})=0, \tag{61}$$

$$\lim_{r\to 0} H(\vec{r})=0, \tag{62}$$

Proof

$$\begin{array}{l}\lim\limits_{r\to 0} r^2E(\vec{r})=0,\\ \lim\limits_{r\to 0} r^2H(\vec{r})=0,\end{array} \tag{63}$$

is derived from (60). the additional condition (60) is reasonable finite energy condition. From GL radial electric integral equation (51)), we have

$$\begin{array}{l} E(\vec{r}')=E^b(\vec{r}')-\\ -\int\limits_{S(r\leq R_2)} \frac{\partial}{\partial r}\left(\left(\left(1-\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}G\right)E\right)dV\\ +\int\limits_{S(r\leq R_2)} \frac{\partial}{\partial r}\left(\left(1-\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}G\right)EdV\\ +\int\limits_{S(r\leq R_2)} \left(\frac{1}{r^2}\left(1-\frac{1}{\varepsilon_r}\right)\right.\\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2G}{\partial\phi^2}\right)\right)EdV\\ +\int\limits_{S(r\leq R_2)} k^2(1-\mu_\theta)GEdV,\end{array} \tag{64}$$

The integral equation (51) is translated to

$$\begin{array}{l} \frac{1}{\varepsilon_\theta}E(\vec{r}')=E^b(\vec{r}')-\\ -\int\limits_{S(r\leq R_2)} \left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{\partial G}{\partial r}EdV\\ +\int\limits_{S(r\leq R_2)} \left(1-\frac{1}{\varepsilon_\theta}\right)\left(\frac{\partial^2}{\partial r^2}G\right)EdV\\ +\int\limits_{S(r\leq R_2)} \left(\frac{1}{r^2}\left(1-\frac{1}{\varepsilon_r}\right)\right.\\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2G}{\partial\phi^2}\right)\right)EdV\\ +\int\limits_{S(r\leq R_2)} k^2(1-\mu_\theta)GEdV,\end{array} \tag{65}$$

The above equation (65) becomes

$$\begin{array}{l} \frac{1}{\varepsilon_\theta}E(\vec{r}')=E^b(\vec{r}')-\\ -\int\limits_{S(r\leq R_2)} \left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{\partial G}{\partial r}EdV\\ -\int\limits_{S(r\leq R_2)} \left(\frac{1}{r^2}\left(1-\frac{1}{\varepsilon_\theta}\right)\right.\\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2G}{\partial\phi^2}\right)\right)EdV\\ -\int\limits_{S(r\leq R_2)} k^2\left(1-\frac{1}{\varepsilon_\theta}\right)GEdV\\ +\int\limits_{S(r\leq R_2)} \left(\frac{1}{r^2}\left(1-\frac{1}{\varepsilon_r}\right)\right.\\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2G}{\partial\phi^2}\right)\right)EdV\\ +\int\limits_{S(r\leq R_2)} k^2(1-\mu_\theta)GEdV.\end{array} \tag{66}$$

The GL electric integral equation (51) becomes to

$$\begin{array}{l} \frac{1}{\varepsilon_\theta}E(\vec{r}')=E^b(\vec{r}')-\\ -\int\limits_{S(r\leq R_2)} \left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{\partial G}{\partial r}EdV\\ +\int\limits_{S(r\leq R_2)} \left(\frac{1}{r^2}\left(\frac{1}{\varepsilon_\theta}-\frac{1}{\varepsilon_r}\right)\right.\\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2G}{\partial\phi^2}\right)\right)EdV\\ +\int\limits_{S(r\leq R_2)} k^2\left(\frac{1}{\varepsilon_\theta}-\mu_\theta\right)GEdV,\end{array} \tag{67}$$

Let

$$G_L(r,r')=G(\vec{r},\vec{r}')-G_0(r,r') \tag{68}$$

to substitute (68) for $G(\vec{r},\vec{r}')$ into (67) , and by using $E_0(r)=0$ in (30), the integral equation (67) becomes

$$\begin{array}{l} \frac{1}{\varepsilon_\theta}E(\vec{r}')=E^b(\vec{r}')-\\ -\int\limits_{S(r\leq R_2)} \left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{\partial G_L}{\partial r}EdV\\ +\int\limits_{S(r\leq R_2)} \left(\frac{1}{r^2}\left(\frac{1}{\varepsilon_\theta}-\frac{1}{\varepsilon_r}\right)\right.\\ \left.\left(\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial G_L}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2G_L}{\partial\phi^2}\right)\right)EdV\\ +\int\limits_{S(r\leq R_2)} k^2\left(\frac{1}{\varepsilon_\theta}-\mu_\theta\right)G_LEdV,\end{array} \tag{69}$$

where

$$\begin{array}{l} G_L(r,r')=G(\vec{r},\vec{r}')-G_0(r,r')\\ =rr'\sum\limits_{l=1}^{\infty} g_l(r,r')\sum\limits_{m=-l}^{l} Y_l^{m*}(\theta,\phi)Y_l^m(\theta',\phi'),\end{array} \tag{70}$$

$$\begin{array}{l} g_l(r,r')\\ =ikj_l(kr)(j_l(kr')-iy_l(kr')), r\leq r'\end{array} \tag{71}$$

$$\begin{array}{l} g_l(r,r')\\ =ik(j_l(kr)-iy_l(kr))j_l(kr'), r\geq r'\end{array} \tag{72}$$

Substitute the $G_L(r,r')$ in (70-72) into the integral equation (69), the equation (69) becomes to the following integral equation,

$$\begin{aligned}
&\tfrac{1}{\varepsilon_\theta}E(\vec{r}') = E^b(\vec{r}') \\
&-r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)\tfrac{\partial}{\partial r}rg_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') \\
&+r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}\tfrac{1}{r^2}\left(\tfrac{1}{\varepsilon_\theta}-\tfrac{1}{\varepsilon_r}\right)l(l+1)rg_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') \\
&+r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}k^2\left(\tfrac{1}{\varepsilon_\theta}-\mu_\theta\right)rg_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi'),
\end{aligned} \tag{73}$$

$$\begin{aligned}
&\lim_{r'\to0}\tfrac{1}{\varepsilon_\theta}E(\vec{r}') = \lim_{r'\to0}E^b(\vec{r}') \\
&-\lim_{r'\to0}r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)\tfrac{\partial}{\partial r}rg_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') \\
&+\lim_{r'\to0}r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}\tfrac{1}{r^2}\left(\tfrac{1}{\varepsilon_\theta}-\tfrac{1}{\varepsilon_r}\right)l(l+1) \\
&rg_l(r,r')\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') \\
&+\lim_{r'\to0}r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}k^2\tfrac{1}{\varepsilon_\theta}rg_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') \\
&-\lim_{r'\to0}r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}k^2\mu_\theta rg_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') \\
&=\lim_{r'\to0}I+\lim_{r'\to0}II+\lim_{r'\to0}III \\
&+\lim_{r'\to0}IV+\lim_{r'\to0}V,
\end{aligned} \tag{74}$$

From the equation (60)

$$\lim_{r'\to0}\frac{1}{\varepsilon_\theta}E(\vec{r}') = 0 \tag{75}$$

From the equation (21)

$$\lim_{r'\to0}I = \lim_{r'\to0}E^b(\vec{r}') = 0, \tag{76}$$

In next, we will prove

$$\begin{aligned}
&\lim_{r'\to0}II = \\
&=-\lim_{r'\to0}r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)\tfrac{\partial}{\partial r}rg_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') = 0,
\end{aligned} \tag{77}$$

$$\begin{aligned}
&\lim_{r'\to0}III = \\
&=\lim_{r'\to0}r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}\left(\tfrac{1}{\varepsilon_\theta}-\tfrac{1}{\varepsilon_r}\right)\tfrac{l(l+1)}{r}g_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') = 0,
\end{aligned} \tag{78}$$

$$\begin{aligned}
&\lim_{r'\to0}IV = \\
&=\lim_{r'\to0}r'\sum_{l=1}^{\infty}\int_{S(r\le R_2)}k^2\tfrac{1}{\varepsilon_\theta}rg_l(r,r') \\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi') = 0,
\end{aligned} \tag{79}$$

Substitute (70) and (71) into the (77) and using LHOPITAL ROLE, we prove the limitation equation (77) in detail,

$$\begin{aligned}
&\lim_{r'\to0}II = \\
&-\lim_{r'\to0}k\sum_{l=1}^{\infty}r'y_l(kr')\int_0^{r'}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)\tfrac{\partial}{\partial r}rj_l(kr)dr \\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to0}k\sum_{l=1}^{\infty}r'j_l(kr')\int_{r'}^{R_2}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)\tfrac{\partial}{\partial r}ry_l(kr)dr \\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to0}ik\sum_{l=1}^{\infty}r'j_l(kr')\int_0^{R_2}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)\tfrac{\partial}{\partial r}rj_l(kr)dr \\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')
\end{aligned} \tag{80}$$

$$\begin{aligned}
&\lim_{r'\to0}II = -\lim_{r'\to0}k\sum_{l=1}^{\infty}r'y_l(kr')\int_0^{r'}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)(l+1)j_l(kr)dr \\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&+\lim_{r'\to0}k\sum_{l=1}^{\infty}r'y_l(kr')\int_0^{r'}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)krj_{l+1}(kr)dr \\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to0}k\sum_{l=1}^{\infty}r'j_l(kr')\int_{r'}^{R_2}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)(l+1)y_l(kr)dr \\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&+\lim_{r'\to0}k\sum_{l=1}^{\infty}r'j_l(kr')\int_{r'}^{R_2}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)kry_{l+1}(kr)dr \\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to0}ik\sum_{l=1}^{\infty}r'j_l(kr')\int_{r'}^{R_2}\left(\tfrac{\partial}{\partial r}\tfrac{1}{\varepsilon_\theta}\right)\tfrac{\partial}{\partial r}rj_l(kr)dr \\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')
\end{aligned} \tag{81}$$

By LHOPITAL ROLE

$$
\begin{aligned}
&\lim_{r'\to 0} II = \\
&\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{(2l)!}{2^l l!}\frac{1}{kl(kr')^{l-1}}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)(l+1)\frac{2^l l!(kr')^l}{(2l+1)!}\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{(2l)!}{2^l l!}\frac{1}{kl(kr')^{l-1}}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{2^{l+1}(l+1)!}{(2l+3)!}(kr')^{l+2}\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{2^l l!}{(2l+1)!}\frac{(kr')^{l+2}}{k(l+1)}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)l\frac{(2l)!}{2^l l!}\frac{1}{(kr')^{l+1}}\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}ik\sum_{l=1}^{\infty}r' j_l(kr')\int_0^{R_2}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}rj_l(kr)dr\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi').
\end{aligned}
\tag{82}
$$

$$
\begin{aligned}
&\lim_{r'\to 0} II = \\
&\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{l+1}{(2l+1)}\frac{r'}{l}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{(kr')^2}{kl(2l+3)(2l+1)}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{l}{(2l+1)}\frac{r'}{(l+1)}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}ik\sum_{l=1}^{\infty}r' j_l(kr')\int_0^{R_2}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}rj_l(kr)dr\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi'),
\end{aligned}
\tag{83}
$$

Because the finite energy condition (60) and GLHUA pre cloak material condition (6.1) *to* (6.4), the integral in above equation (83) is finite integrative,

$$
\begin{aligned}
&\lim_{r'\to 0} II = \\
&\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{l+1}{(2l+1)}\frac{r'}{l}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{(kr')^2}{kl(2l+3)(2l+1)}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}\sum_{l=1}^{\infty}\frac{l}{(2l+1)}\frac{r'}{(l+1)}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&-\lim_{r'\to 0}ik\sum_{l=1}^{\infty}r' j_l(kr')\int_0^{R_2}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}rj_l(kr)dr\\
&\sum_{m=-l}^{l}\int_0^{\pi}\int_0^{2\pi}Y_l^{m*}(\theta,\phi)E(r,\theta,\phi)\sin\theta d\theta d\phi Y_l^m(\theta',\phi')\\
&=0,
\end{aligned}
\tag{84}
$$

Therefore,

$$
\begin{aligned}
&\lim_{r'\to 0} II = -\lim_{r'\to 0} r'\sum_{l=1}^{\infty}\int\limits_{S(r\le R_2)}\left(\frac{\partial}{\partial r}\frac{1}{\varepsilon_\theta}\right)\frac{\partial}{\partial r}rg_l(r,r')\\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi')=0,
\end{aligned}
\tag{85}
$$

limitation equation (77) is proved. Similarly, we can prove limitation equation (78)

$$
\begin{aligned}
&\lim_{r'\to 0} III = \lim_{r'\to 0} r'\sum_{l=1}^{\infty}\int\limits_{S(r\le R_2)}\frac{1}{r^2}\left(\frac{1}{\varepsilon_\theta}-\frac{1}{\varepsilon_r}\right)l(l+1)rg_l(r,r')\\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi')=0,
\end{aligned}
\tag{86}
$$

, and (79),

$$
\begin{aligned}
&\lim_{r'\to 0} IV = \lim_{r'\to 0} r'\sum_{l=1}^{\infty}\int\limits_{S(r\le R_2)}k^2\frac{1}{\varepsilon_\theta}rg_l(r,r')\\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi')=0,
\end{aligned}
\tag{87}
$$

Substitute (75)-(79) into the limitation equation (74), the limitation equation (74) induces

$$
\begin{aligned}
&\lim_{r'\to 0} V = -\lim_{r'\to 0} r'\sum_{l=1}^{\infty}\int\limits_{S(r\le R_2)}k^2\mu_\theta rg_l(r,r')\\
&\sum_{m=-l}^{l}Y_l^{m*}(\theta,\phi)EdVY_l^m(\theta',\phi')=0,
\end{aligned}
\tag{88}
$$

Substitute (71) and (72) for $g_l(r,r')$ into the (88),

$$\begin{aligned}
&\lim_{r'\to 0} V = \\
&-\lim_{r'\to 0} k^3 \sum_{l=1}^{\infty} r' y_l(kr') \int_0^{r'} \mu_\theta r j_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to 0} k^3 \sum_{l=1}^{\infty} r' j_l(kr') \int_{r'}^{R_2} \mu_\theta r y_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to 0} ik^3 \sum_{l=1}^{\infty} r' j_l(kr') \int_0^{R_2} \mu_\theta r j_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&= 0
\end{aligned} \tag{89}$$

$$\begin{aligned}
&\lim_{r'\to 0} V = \\
&+\lim_{r'\to 0} k^2 \sum_{l=1}^{\infty} \tfrac{(2l)!}{2^l l!} \tfrac{1}{(kr')^l} \int_0^{r'} \mu_\theta r j_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to 0} k^2 \sum_{l=1}^{\infty} \tfrac{2^l l! (kr')^{l+1}}{(2l+1)!} \int_{r'}^{R_2} \mu_\theta r y_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to 0} ik^3 \sum_{l=1}^{\infty} r' j_l(kr') \int_0^{R_2} \mu_\theta r j_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&= 0,
\end{aligned} \tag{90}$$

By LHOPITAL ROLE

$$\begin{aligned}
&\lim_{r'\to 0} V = \lim_{r'\to 0} k^2 \sum_{l=1}^{\infty} \tfrac{(2l)!}{2^l l!} \tfrac{1}{kl(kr')^{l-1}} \mu_\theta r' j_l(kr') \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to 0} k^2 \sum_{l=1}^{\infty} \tfrac{2^l l! (kr')^{l+2}}{k(l+1)(2l+1)!} \mu_\theta r' y_l(kr') \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to 0} ik^3 \sum_{l=1}^{\infty} r' j_l(kr') \int_0^{R_2} \mu_\theta r j_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&= 0,
\end{aligned} \tag{91}$$

$$\begin{aligned}
&\lim_{r'\to 0} V = \\
&\lim_{r'\to 0} k^2 \sum_{l=1}^{\infty} \tfrac{(2l)!}{2^l l!} \tfrac{1}{kl(kr')^{l-1}} \tfrac{f(r')}{r'} \tfrac{2^l l!}{(2l+1)!} (kr')^l \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&+\lim_{r'\to 0} k^2 \sum_{l=1}^{\infty} \tfrac{2^l l! (kr')^{l+2}}{k(l+1)(2l+1)!} \tfrac{f(r')}{r'} \tfrac{(2l)!}{2^l l!} \tfrac{1}{(kr')^{l+1}} \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to 0} ik^3 \sum_{l=1}^{\infty} r' j_l(kr') \int_0^{R_2} \varepsilon_\theta r j_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&= 0,
\end{aligned} \tag{92}$$

$$\begin{aligned}
&\lim_{r'\to 0} V = \\
&\lim_{r'\to 0} k^2 f(r') \sum_{l=1}^{\infty} \tfrac{1}{l(2l+1)} \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&+\lim_{r'\to 0} k^2 f(r') \sum_{l=1}^{\infty} \tfrac{1}{(l+1)(2l+1)} \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&-\lim_{r'\to 0} ik^3 \sum_{l=1}^{\infty} r' j_l(kr') \int_0^{R_2} \varepsilon_\theta r j_l(kr) dr \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi Y_l^m(\theta',\phi') \\
&= 0,
\end{aligned} \tag{93}$$

$$\begin{aligned}
&\lim_{r'\to 0} k^2 R_2^2 \sum_{l=1}^{\infty} \tfrac{1}{l(l+1)} \\
&\sum_{m=-l}^{l} \int_0^{\pi} \int_0^{2\pi} Y_l^{m*}(\theta,\phi) E(r,\theta,\phi) \sin\theta d\theta d\phi \\
&Y_l^m(\theta',\phi') = 0,
\end{aligned} \tag{94}$$

Because for incident wave with electric current point source $\delta(\vec{r}-\vec{r}_s)\vec{e}_s$ at (r_s,θ_s,ϕ_s) $r_s > R_O > R_2$, $E_0(r) = 0$ in (30),

$$\begin{aligned}
&E(r',\theta',\phi') = \sum_{l=1}^{\infty} E_l(r') \\
&\sum_{m=-l}^{l} Y_l^m(\theta',\phi') Y_l^m(\theta_s,\phi_s)
\end{aligned} \tag{95}$$

Substitute (95) into the (94), we have

$$\begin{aligned}
&\lim_{r'\to 0} k^2 R_2^2 \sum_{l=1}^{\infty} \tfrac{1}{l(l+1)} E_l(r') \\
&\sum_{m=-l}^{l} Y_l^m(\theta',\phi') Y_l^m(\theta_s,\phi_s) = 0,
\end{aligned} \tag{96}$$

Let

$$\Re(\theta,\phi)=\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial}{\partial\theta}+\frac{1}{\sin^2\theta}\frac{\partial^2}{\partial\phi^2}, \tag{97}$$

To use $\Re(\theta,\phi)$ in (97) to make action to both side of the limitation equation (96), we have

$$\begin{array}{l}\lim\limits_{r'\to 0}k^2R_2^2\Re(\theta',\phi')\sum\limits_{l=1}^{\infty}\frac{1}{l(l+1)}E_l(r')\\ \sum\limits_{m=-l}^{l}Y_l^m(\theta',\phi')Y_l^m(\theta_s,\phi_s)=0\end{array} \tag{98}$$

$$\lim_{r'\to 0}k^2R_2^2E(r',\theta',\phi')=0, \tag{99}$$

For every (θ',ϕ') and $k\geq k_0>0$

$$\lim_{r'\to 0}E(r',\theta',\phi')=0, \tag{100}$$

Other proof approach is as follows: By using above similar with proof process of (98), for every term l and k, we can prove that

$$\lim_{r'\to 0}k^2R_2^2\frac{1}{l(l+1)}E_l(r')=0, \tag{101}$$

, Because $E_0(r)=0$ in (30),$l\geq 1$ and for $k\geq k_0>0$

$$\lim_{r'\to 0}E_l(r')=0, \tag{102}$$

$$\begin{array}{l}\lim\limits_{r'\to 0}E(r',\theta',\phi')=\sum\limits_{l=1}^{\infty}\lim\limits_{r'\to 0}E_l(r')\\ \sum\limits_{m=-l}^{l}Y_l^m(\theta',\phi')Y_l^m(\theta_s,\phi_s)=0,\end{array} \tag{103}$$

The limitation equation (61) is proved. Similarly, we can prove limitation equation (62)

$$\begin{array}{l}\lim\limits_{r'\to 0}H(r',\theta',\phi')=\sum\limits_{l=1}^{\infty}\lim\limits_{r'\to 0}H_l(r')\\ \sum\limits_{m=-l}^{l}Y_l^m(\theta',\phi')Y_l^m(\theta_s,\phi_s)=0,\end{array} \tag{104}$$

Similarly, for incident plane electromagnetic wave , we also can prove limitation equation (61) and (62). The theorem 6.1 is proved.

***Theorem* 6.2**, Suppose that the anisotropic relative electric permittivity $\varepsilon_r(r)$,$\varepsilon_\theta(r)$ $\varepsilon_\phi(r)$ and magnetic permeability, $\mu_r(r)$,$\mu_\theta(r)$ $\mu_\phi(r)$, satisfy the above GL-HUA pre cloak material conditions in invisible sphere (6.1) *to* (6.4), and finite energy condition (60), then

$$\lim_{r\to 0}\frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}E(\vec{r})=0, \tag{105}$$

$$\lim_{r\to 0}\frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}H(\vec{r})=0, \tag{106}$$

***Proof* :**, Using similar proof process, the theorem 6.2 can be proved.

***Theorem* 6.3**, Suppose that the anisotropic relative electric permittivity $\varepsilon_r(r)$,$\varepsilon_\theta(r)$ $\varepsilon_\phi(r)$ and magnetic permeability, $\mu_r(r)$,$\mu_\theta(r)$ $\mu_\phi(r)$, satisfy the above GL-HUA pre cloak material conditions in invisible sphere (6.1) *to* (6.4), also angular electromagnetic wave satisfy following finite energy condition (63)

$$\begin{array}{l}\int\limits_{S(r\leq R_2)}\Big(|rE_\theta(\vec{r})|^2+|rH_\theta(\vec{r})|^2\\ +|rE_\phi(\vec{r})|^2+|rH_\phi(\vec{r})|^2\Big)\,dV,\\ is\ finite,\end{array} \tag{107}$$

then

$$\begin{array}{l}\lim\limits_{r\to 0}rE_\theta(\vec{r})=0,\\ \lim\limits_{r\to 0}rE_\phi(\vec{r})=0,\end{array} \tag{108}$$

$$\begin{array}{l}\lim\limits_{r\to 0}rH_\theta(\vec{r})=0,\\ \lim\limits_{r\to 0}rH_\phi(\vec{r})=0,\end{array} \tag{109}$$

***Proof* :**, Because the source is outside sphere,$r_s>R_O>R_2$, In the Sphere $0<r\leq R_2$, from (4), we have

$$\begin{array}{l}\nabla\cdot\vec{D}=\frac{1}{r^2}\frac{\partial}{\partial r}\left(r^2\varepsilon_rE_r\right)+\\ +\frac{1}{\sin\theta r}\frac{\partial}{\partial\theta}\sin\theta\varepsilon_\theta E_\theta+\frac{1}{r\sin\theta}\frac{\partial}{\partial\phi}\varepsilon_\phi E_\phi=0\end{array} \tag{110}$$

The $\vec{D}$ is displacement electric in spherical coordinate, by definition of GL electromagnetic wave (18), equation (110) becomes

$$\begin{array}{l}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta rE_\theta+\frac{1}{\sin\theta}\frac{\partial}{\partial\phi}rE_\phi\\ =-\frac{1}{\varepsilon_\theta}\frac{\partial E}{\partial r},\end{array} \tag{111}$$

By Maxwell equation (1) and (18), we have

$$\begin{array}{l}-\frac{1}{\sin\theta}\frac{\partial}{\partial\phi}rE_\theta+\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta rE_\phi\\ =-i\omega\mu_0H,\end{array} \tag{112}$$

Rewrite (111) and (112) as matrix equation

$$\begin{aligned}&\left[\begin{array}{cc}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta & \frac{1}{\sin\theta}\frac{\partial}{\partial\phi}\\ -\frac{1}{\sin\theta}\frac{\partial}{\partial\phi} & \frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\end{array}\right]\left[\begin{array}{c}rE_\theta\\ rE_\phi\end{array}\right]=\\&=\left[\begin{array}{c}-\frac{1}{\varepsilon_\theta}\frac{\partial E}{\partial r}\\ -i\omega\mu_0 H\end{array}\right]\end{aligned}\tag{113}$$

The adjoint Greens equation of equation (113) on $[0,\pi;0,2\pi]$ is

$$\begin{aligned}&\left[\begin{array}{cc}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta & -\frac{1}{\sin\theta}\frac{\partial}{\partial\phi}\\ \frac{1}{\sin\theta}\frac{\partial}{\partial\phi} & \frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\end{array}\right]\\&\left[\begin{array}{cc}G_{11}(\theta,\theta',\phi,\phi') & G_{21}(\theta,\theta',\phi,\phi')\\ G_{12}(\theta,\theta',\phi,\phi') & G_{22}(\theta,\theta',\phi,\phi')\end{array}\right]\\&=\frac{1}{\sin\theta}\left[\begin{array}{cc}\delta(\theta,\theta',\phi,\phi') & \\ & \delta(\theta,\theta',\phi,\phi')\end{array}\right],\end{aligned}\tag{114}$$

It is different from GL Green equation in (44), the adjoint Greens equation is novel GLHUA angular Green equation. It is different from GL Green function in (45), the GLHUA angular Green function matrix is in the following

$$\begin{aligned}&G_{11}(\theta,\phi,\theta',\phi')=\\&\sum_{l=}^{\infty}\frac{1}{l(l+1)}\sum_{m=-l}^{l}\frac{\partial}{\partial\theta}Y_l^{m*}(\theta,\phi)Y_l^{m*}(\theta',\phi'),\\&G_{12}(\theta,\phi,\theta',\phi')=\\&-\sum_{l=}^{\infty}\frac{1}{l(l+1)}\sum_{m=-l}^{l}\frac{\partial}{\sin\theta\partial\phi}Y_l^{m*}(\theta,\phi)Y_l^{m*}(\theta',\phi'),\\&G_{21}(\theta,\phi,\theta',\phi')=\\&\sum_{l=}^{\infty}\frac{1}{l(l+1)}\sum_{m=-l}^{l}\frac{\partial}{\sin\theta\partial\phi}Y_l^{m*}(\theta,\phi)Y_l^{m*}(\theta',\phi'),\\&G_{22}(\theta,\phi,\theta',\phi')=\\&\sum_{l=}^{\infty}\frac{1}{l(l+1)}\sum_{m=-l}^{l}\frac{\partial}{\partial\theta}Y_l^{m*}(\theta,\phi)Y_l^{m*}(\theta',\phi'),\end{aligned}\tag{115}$$

To use product of GLHUA Greens function matrix by $\sin\theta$, $G(\theta,\theta',\phi,\phi')\sin\theta$, to multiply the matrix equation (113) and take sphere surface integral of resulted equation on $[0,\pi;0,2\pi]$

$$\begin{aligned}&\int_0^\pi\int_0^{2\pi}\left(\left[\begin{array}{cc}G_{11} & G_{12}\\ G_{21} & G_{22}\end{array}\right]\right.\\&\left.\left[\begin{array}{cc}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta & \frac{1}{\sin\theta}\frac{\partial}{\partial\phi}\\ -\frac{1}{\sin\theta}\frac{\partial}{\partial\phi} & \frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\end{array}\right]\left[\begin{array}{c}rE_\theta\\ rE_\phi\end{array}\right]\right)\sin\theta d\theta d\phi\\&=\int_0^\pi\int_0^{2\pi}\left[\begin{array}{cc}G_{11} & G_{12}\\ G_{21} & G_{22}\end{array}\right]\left[\begin{array}{c}-\frac{1}{\varepsilon_\theta}\frac{\partial E}{\partial r}\\ -i\omega\mu_0 H\end{array}\right]\sin\theta d\theta d\phi,\end{aligned}\tag{116}$$

To use vector $[rE_\theta, rE_\phi]\sin\theta$ to multiply the adjoint Greens equation of equation (114) and take sphere surface integral of resulted adjoint equation on $[0,\pi;0,2\pi]$,

$$\begin{aligned}&\int_0^\pi\int_0^{2\pi}\left[\begin{array}{cc}G_{11} & G_{12}\\ G_{21} & G_{22}\end{array}\right]\\&\left[\begin{array}{cc}\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta & \frac{1}{\sin\theta}\frac{\partial}{\partial\phi}\\ -\frac{1}{\sin\theta}\frac{\partial}{\partial\phi} & \frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\end{array}\right]\left[\begin{array}{c}rE_\theta\\ rE_\phi\end{array}\right]\sin\theta d\theta d\phi\\&=\int_0^\pi\int_0^{2\pi}\left[\begin{array}{cc}\delta & \\ & \delta\end{array}\right]\left[\begin{array}{c}rE_\theta\\ rE_\phi\end{array}\right]\sin\theta d\theta d\phi,\end{aligned}\tag{117}$$

To subtract equation (116) from (117), we have

$$\begin{aligned}&\left[\begin{array}{c}rE_\theta(r,\theta',\phi')\\ rE_\phi(r,\theta',\phi')\end{array}\right]=\\&=\int_0^\pi\int_0^{2\pi}\left[\begin{array}{cc}G_{11} & G_{12}\\ G_{21} & G_{22}\end{array}\right]\left[\begin{array}{c}-\frac{1}{\varepsilon_\theta}\frac{\partial E}{\partial r}\\ -i\omega\mu_0 H\end{array}\right]\sin\theta d\theta d\phi\end{aligned}\tag{118}$$

Based on theorem 6.1 and theorem 6.2, we have

$$\begin{aligned}&\lim_{r\to0}\left[\begin{array}{c}rE_\theta(r,\theta',\phi')\\ rE_\phi(r,\theta',\phi')\end{array}\right]\\&=\lim_{r\to0}\int_0^\pi\int_0^{2\pi}\left[\begin{array}{cc}G_{11} & G_{12}\\ G_{21} & G_{22}\end{array}\right]\left[\begin{array}{c}-\frac{1}{\varepsilon_\theta}\frac{\partial E}{\partial r}\\ -i\omega\mu_0 H\end{array}\right]\sin\theta d\theta d\phi\\&=\int_0^\pi\int_0^{2\pi}\left[\begin{array}{cc}G_{11} & G_{12}\\ G_{21} & G_{22}\end{array}\right]\lim_{r\to0}\left[\begin{array}{c}-\frac{1}{\varepsilon_\theta}\frac{\partial E}{\partial r}\\ -i\omega\mu_0 H\end{array}\right]\sin\theta d\theta d\phi\\&=\left[\begin{array}{c}0\\ 0\end{array}\right],\end{aligned}\tag{119}$$

We have proved the first part of (108) of theorem 6.3, similarly, we can prove second part of (109). The theorem 6.3 is proved.

***Theorem* 6.4**, Suppose that the anisotropic relative electric permittivity $\varepsilon_r(r)$,$\varepsilon_\theta(r)$ $\varepsilon_\phi(r)$ and magnetic permeability, $\mu_r(r)$,$\mu_\theta(r)$ $\mu_\phi(r)$, satisfy the above GLHUA pre cloak material conditions in invisible sphere (6.1) *to* (6.4), also angular electromagnetic wave satisfy following finite energy condition (63), then

$$\begin{aligned}&\lim_{r\to0}\frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}rE_\theta(\vec{r})=0,\\&\lim_{r\to0}\frac{1}{\varepsilon_\theta}\frac{\partial}{\partial r}rE_\phi(\vec{r})=0,\end{aligned}\tag{120}$$

$$\begin{aligned}&\lim_{r\to0}\frac{1}{\mu_\theta}\frac{\partial}{\partial r}rH_\theta(\vec{r})=0,\\&\lim_{r\to0}\frac{1}{\mu_\theta}\frac{\partial}{\partial r}rH_\phi(\vec{r})=0,\end{aligned}\tag{121}$$

***Proof* :**, By using the similar proof process on the 8.3, we can prove the theorem 6.4.

VII. DISCUSSION AND CONCLUSION

The pre cloak material conditions (6.1) *to* (6.4) in GLHUA sphere is from GL zero scattering inversion and GL no scattering modeling. Many GL no scattering modeling simulations show that under the conditions

(6.1) *to* (6.4), $\lim_{r\to 0} E(r) = 0$,, and $\lim_{r\to 0} H(r) = 0$, are verified. These condition is not unique and can be relaxed. We publish this paper to arXiv for support our paper arXiv.org/abs/1612.02857. Our GLHUA cloak and GLHUA sphere publication in arXiv are for open review. Please colleague give comments to me by my email or give open comments in arXiv. All copyright and patent of the GLHUA EM cloaks,GLHUA sphere and GL modeling and inversion methods are reserved by authors in GL Geophysical Laboratory.If some colleague cite our paper in his work paper, please cite our paper as reference in his paper..

Acknowledgments

We wish to acknowledge the support of the GL Geophysical Laboratory and thank the GLGEO Laboratory to approve the paper publication.

[1] Xie, G., J.H. Li, E. Majer, D. Zuo, M. Oristaglio "3-D electromagnetic modeling and nonlinear inversion," *Geophysics*, Vol. 65, No. 3, 804–822, 2000.

[2] Xie, G., F. Xie, L. Xie, and J. Li, "New GL method and its advantages for resolving historical diffculties," *Progress In Electromagnetics Research*, PIER 63, 141–152, 2006.

[3] Xie, G., J. Li, L. Xie, and F. Xie, "GL metro carlo EM inversion," *Journal of Electromagnetic Waves and Applications*, Vol. 20, No. 14, 1991–2000, 2006.